

ON POSITIVE WEAK SOLUTIONS FOR A CLASS OF WEIGHTED (P,Q)-LAPLACIAN NONLINEAR SYSTEM

SALAH A. KHAFAGY

ABSTRACT. In this paper we study the existence and nonexistence of positive weak solutions for the nonlinear system

$$\left. \begin{aligned} -\Delta_{P,p}u &= \lambda a(x)v^\gamma + b(x)u^\alpha && \text{in } \Omega, \\ -\Delta_{Q,q}v &= \lambda c(x)u^\delta + d(x)v^\beta && \text{in } \Omega, \\ u = v &= 0 && \text{on } \partial\Omega. \end{aligned} \right\}$$

where $\Delta_{R,r}$ with $r > 1$ and $R = R(x)$ is a weight function, denotes the weighted r -Laplacian defined by $\Delta_{R,r}u \equiv \operatorname{div}[R(x)|\nabla u|^{r-2}\nabla u]$, λ is a positive parameter, $a(x)$, $b(x)$, $c(x)$ and $d(x)$ are weight functions, $\alpha, \beta \geq 0$, $\gamma, \delta > 0$, $\alpha + \delta < p - 1$, $\beta + \gamma < q - 1$ and $\Omega \subset \mathbb{R}^N$ is a bounded domain with smooth boundary $\partial\Omega$. We use the method of sub-supersolutions to establish our results.

Mathematics Subject Classification (2010): 35D30, 93C10

Key words: Weak solution, Nonlinear system, Sub-supersolutions method

Article history:

Received 27 November 2016

Received in revised form 12 August 2017

Accepted 23 September 2017

1. INTRODUCTION

Many authors are interested in the study of the existence, nonexistence and stability of positive weak solutions of nonlinear systems involving weighted p -Laplacian during recent years (see [2, 4, 5, 12, 13, 20] and the references therein).

Khafagy in [15] proved the existence, nonexistence and uniqueness of positive weak solution for the nonlinear system

$$(1.1) \quad \left\{ \begin{aligned} -\Delta_{P,p}u &= \lambda a(x)u^\alpha && \text{in } \Omega \\ u &> k && \text{in } \Omega \\ u &= k && \text{on } \partial\Omega, \end{aligned} \right.$$

where $0 < \alpha < p - 1$, using the method of sub-supersolutions. These results extended in [8] to the nonlinear system

$$(1.2) \quad \left\{ \begin{aligned} -\Delta_{P,p}u &= a(x)(\lambda u^\alpha + u^\beta) && \text{in } \Omega \\ u &> k && \text{in } \Omega \\ u &= k && \text{on } \partial\Omega, \end{aligned} \right.$$

where $0 < \beta \leq \alpha < p - 1$.

In [7] the author studied the existence and nonexistence of positive weak solutions for the weighted (p, q) -Laplacian nonlinear system

$$(1.3) \quad \left. \begin{aligned} -\Delta_{P,p}u &= \lambda a(x)v^\beta & \text{in } \Omega, \\ -\Delta_{Q,q}v &= \lambda b(x)u^\alpha & \text{in } \Omega, \\ u = v &= 0 & \text{on } \partial\Omega, \end{aligned} \right\}$$

where $0 < \alpha < p - 1$, $0 < \beta < q - 1$, using the the sub-supersolutions method.

In this paper, we are concerned with the existence and nonexistence of positive weak solutions for the nonlinear system

$$(1.4) \quad \left. \begin{aligned} -\Delta_{P,p}u &= \lambda a(x)v^\gamma + b(x)u^\alpha & \text{in } \Omega, \\ -\Delta_{Q,q}v &= \lambda c(x)u^\delta + d(x)v^\beta & \text{in } \Omega, \\ u = v &= 0 & \text{on } \partial\Omega, \end{aligned} \right\}$$

where $\Delta_{R,r}$ with $r > 1$ and $R = R(x)$ is a weight function, $R(x) = P(x)$ when $r = p$ and $R(x) = Q(x)$ when $r = q$, denotes the weighted r -Laplacian defined by $\Delta_{R,r}u \equiv \operatorname{div}[R(x)|\nabla u|^{r-2}\nabla u]$, λ is a positive parameter, $a(x)$, $b(x)$, $c(x)$ and $d(x)$ are weight functions, $\alpha, \beta \geq 0$, $\gamma, \delta > 0$, $\alpha + \delta < p - 1$, $\beta + \gamma < q - 1$ and $\Omega \subset \mathbb{R}^N$ is a bounded domain with smooth boundary $\partial\Omega$. We discuss, under some certian conditions, the existence of positive weak solution for system (1.4). We also study the conditions under which system (1.4) has no positive weak solution. The existence results are obtained by the sub-supersolutions method.

Clearly, our work extends the work in (1.1)-(1.3) by considering a problem with a more general class of functions. Therefore we have a generalization of all results obtained for the above systems.

On the other hand, the existence of weak solutions for nonlinear systems involving p -Laplacian operators with different weights have been studied using, the sub-supersolutions method (see [8, 10, 11, 14, 15]) the theory of nonlinear monotone operators method (see [16, 21]) and the approximation method (see [22, 23]).

Finally, let us explain the plan of the paper. In section 2, we introduce some technical results and notations, which are established in [6]. In section 3, we prove the existence of a positive weak solutions for system (1.4) by using the method of sub-supersolutions. In section 4, we consider the nonexistence results.

2. TECHNICAL RESULTS

In this section, we introduce some technical results [6] concerning the degenerated homogeneous eigenvalue problem

$$(2.1) \quad \left. \begin{aligned} -\Delta_{R,r}u &= -\operatorname{div}[R(x)|\nabla u|^{r-2}\nabla u] = \lambda S(x)|u|^{r-2}u & \text{in } \Omega, \\ u &= 0 & \text{on } \partial\Omega, \end{aligned} \right\}$$

where $R(x)$ is a weight function (measurable and positive a.e. in Ω), satisfying the conditions

$$(2.2) \quad R(x) \in L^1_{Loc}(\Omega), \quad (R(x))^{-\frac{1}{r-1}} \in L^1_{Loc}(\Omega), \quad \text{with } r > 1,$$

$$(2.3) \quad (R(x))^{-s} \in L^1(\Omega), \quad \text{with } s \in \left(\frac{N}{r}, \infty\right) \cap \left[\frac{1}{r-1}, \infty\right),$$

and $S(x)$ is a measurable function satisfies

$$(2.4) \quad S(x) \in L^{\frac{k}{k-r}}(\Omega),$$

with some k satisfies $r < k < r_s^*$ where $r_s^* = \frac{Nr_s}{N-r_s}$ with $r_s = \frac{rs}{s+1} < r < r_s^*$ and $\operatorname{meas} \{x \in \Omega : S(x) > 0\} > 0$.

Lemma 2.1. [6] *There exists the least(i.e. the first or principal) eigenvalue $\lambda_1^{(r)} > 0$ and precisely one corresponding eigenfunction $\phi_{1,r} \geq 0$ a.e. in Ω ($\phi_{1,r}$ not identical to 0) of the eigenvalue problem (2.1).*

Moreover, it is characterized by

$$(2.5) \quad \lambda_1^{(r)} \int_{\Omega} S(x) \phi_{1,r}^r \leq \int_{\Omega} R(x) |\nabla \phi_{1,r}^r|^p.$$

Lemma 2.2. [6] *Let $\phi_{1,r} \geq 0$ a.e. in Ω , be the eigenfunction corresponding to the first eigenvalue $\lambda_1^{(r)} > 0$ of the eigenvalue problem (2.1). Then $\phi_{1,r} \in L^\infty(\Omega)$.*

Now, let us introduce the weighted Sobolev space $W^{1,p}(R, \Omega)$ which is the set of all real valued functions u defined in Ω for which (see [6])

$$(2.6) \quad \|u\|_{W^{1,r}(R,\Omega)} = \left[\int_{\Omega} |u|^r + \int_{\Omega} R(x) |\nabla u|^r \right]^{\frac{1}{r}} < \infty.$$

Since we are dealing with the Dirichlet problem, we introduce also the space $W_0^{1,r}(R, \Omega)$ as the closure of $C_0^\infty(\Omega)$ in $W^{1,r}(R, \Omega)$ with respect to the norm

$$(2.7) \quad \|u\|_{W_0^{1,r}(R,\Omega)} = \left[\int_{\Omega} R(x) |\nabla u|^r \right]^{\frac{1}{r}} < \infty,$$

which is equivalent to the norm given by (2.6). Both spaces $W^{1,r}(R, \Omega)$ and $W_0^{1,r}(R, \Omega)$ are well defined reflexive Banach Spaces.

3. EXISTENCE RESULTS

We shall prove the existence of a positive weak solution for system (1.4) by constructing a subsolution $(\psi_1, \psi_2) \in W_0^{1,p}(P, \Omega) \times W_0^{1,q}(Q, \Omega)$ and a supersolution $(z_1, z_2) \in W_0^{1,p}(P, \Omega) \times W_0^{1,q}(Q, \Omega)$ of (1.4) such that $\psi_i \leq z_i$ for $i = 1, 2$. That is, $\psi_i, i = 1, 2$, satisfies

$$\begin{aligned} \int_{\Omega} P(x) |\nabla \psi_1|^{p-2} \nabla \psi_1 \nabla \zeta dx &\leq \lambda \int_{\Omega} a(x) \psi_2^\gamma \zeta dx + \int_{\Omega} b(x) \psi_1^\alpha \zeta dx, \\ \int_{\Omega} Q(x) |\nabla \psi_2|^{q-2} \nabla \psi_2 \nabla \eta dx &\leq \lambda \int_{\Omega} c(x) \psi_1^\delta \eta dx + \int_{\Omega} d(x) \psi_2^\beta \eta dx, \end{aligned}$$

and $z_i, i = 1, 2$, satisfies

$$\begin{aligned} \int_{\Omega} P(x) |\nabla z_1|^{p-2} \nabla z_1 \nabla \zeta dx &\geq \lambda \int_{\Omega} a(x) z_2^\gamma \zeta dx + \int_{\Omega} b(x) z_1^\alpha \zeta dx, \\ \int_{\Omega} Q(x) |\nabla z_2|^{q-2} \nabla z_2 \nabla \eta dx &\geq \lambda \int_{\Omega} c(x) z_1^\delta \eta dx + \int_{\Omega} d(x) z_2^\beta \eta dx, \end{aligned}$$

for all test functions $\zeta \in W_0^{1,p}(P, \Omega)$ and $\eta \in W_0^{1,q}(Q, \Omega)$ with $\zeta, \eta \geq 0$.

Lemma 3.1. (see [3, 7]) *Suppose there exist sub and supersolutions (ψ_1, ψ_2) and (z_1, z_2) respectively of (1.4) such that $(\psi_1, \psi_2) \leq (z_1, z_2)$. Then (1.4) has a solution (u, v) such that $(u, v) \in [(\psi_1, \psi_2), (z_1, z_2)]$.*

Theorem 3.2. *Let $(p-1)(q-1) - \gamma\delta > 0$. Then system (1.4) has a positive weak solution (u, v) for each $\lambda > 0$.*

Proof. Let $\lambda_1^{(r)}$ be the first eigenvalue of the eigenvalue problem (2.1) and $\phi_{1,r}$ the corresponding positive eigenfunction satisfying $\phi_{1,r} > 0$ in Ω and $|\nabla\phi_{1,r}| > 0$ on $\partial\Omega$ with $\|\phi_{1,r}\|_\infty = 1$, for $r = p, q$. Then we have

$$(3.1) \quad \left. \begin{aligned} -\Delta_{P,p}\phi_{1,p} &= \lambda_1^{(p)}a(x)|\phi_{1,p}|^{p-2}\phi_{1,p} & \text{in } \Omega, & \quad \phi_{1,p} = 0 & \text{on } \partial\Omega, \\ -\Delta_{Q,q}\phi_{1,q} &= \lambda_1^{(q)}c(x)|\phi_{1,q}|^{q-2}\phi_{1,q} & \text{in } \Omega, & \quad \phi_{1,q} = 0 & \text{on } \partial\Omega. \end{aligned} \right\}$$

Since $(p-1)(q-1) - \gamma\delta > 0$, we can take k such that

$$(3.2) \quad \frac{\delta}{q-1} < k < \frac{p-1}{\gamma}.$$

We shall verify that $(\psi_1, \psi_2) = (\xi\phi_{1,p}^{\frac{p}{p-1}}, \xi^k\phi_{1,q}^{\frac{q}{q-1}})$ is a subsolution of (1.4), where $\xi > 0$ is small and specified later. Let $\zeta \in W_0^{1,p}(P, \Omega)$ with $\zeta \geq 0$. A calculation shows that

$$\begin{aligned} \int_{\Omega} P(x)|\nabla\psi_1|^{p-2}\nabla\psi_1 \cdot \nabla\zeta dx &= \left(\frac{p}{p-1}\xi\right)^{p-1} \int_{\Omega} P(x)\phi_{1,p}|\nabla\phi_{1,p}|^{p-2}\nabla\phi_{1,p} \cdot \nabla\zeta dx \\ &= \left(\frac{p}{p-1}\xi\right)^{p-1} \int_{\Omega} (P(x)|\nabla\phi_{1,p}|^{p-2}\nabla\phi_{1,p}\nabla(\phi_{1,p}\zeta) - P(x)|\nabla\phi_{1,p}|^p)\zeta dx \\ &= \left(\frac{p}{p-1}\xi\right)^{p-1} \int_{\Omega} (\lambda_1^{(p)}a(x)\phi_{1,p}^p - P(x)|\nabla\phi_{1,p}|^p)\zeta dx. \end{aligned}$$

Similarly, for $\eta \in W_0^{1,q}(Q, \Omega)$ with $\eta \geq 0$, we have

$$\int_{\Omega} Q(x)|\nabla\psi_2|^{q-2}\nabla\psi_2 \cdot \nabla\eta dx = \left(\frac{q}{q-1}\xi^k\right)^{q-1} \int_{\Omega} (\lambda_1^{(q)}c(x)\phi_{1,q}^q - Q(x)|\nabla\phi_{1,q}|^q)\eta dx.$$

Since $\phi_{1,r} = 0$ and $|\nabla\phi_{1,r}| > 0$ on $\partial\Omega$, there is $\epsilon > 0$ such that

$$\lambda_1^{(p)}a(x)\phi_{1,p}^p - P(x)|\nabla\phi_{1,p}|^p \leq 0 \quad \text{and} \quad \lambda_1^{(q)}c(x)\phi_{1,q}^q - Q(x)|\nabla\phi_{1,q}|^q \leq 0 \quad \text{on } \bar{\Omega}_\epsilon$$

with $\bar{\Omega}_\epsilon = \{x \in \Omega : d(x, \partial\Omega) \leq \epsilon\}$. This shows that

$$\left(\frac{p}{p-1}\xi\right)^{p-1} \int_{\bar{\Omega}_\epsilon} (\lambda_1^{(p)}a(x)\phi_{1,p}^p - P(x)|\nabla\phi_{1,p}|^p)\zeta dx \leq 0 \leq \lambda \int_{\bar{\Omega}_\epsilon} a(x)\psi_2^\gamma \zeta dx + \int_{\bar{\Omega}_\epsilon} b(x)\psi_1^\alpha \zeta dx,$$

and

$$\left(\frac{q}{q-1}\xi^k\right)^{q-1} \int_{\bar{\Omega}_\epsilon} (\lambda_1^{(q)}c(x)\phi_{1,q}^q - Q(x)|\nabla\phi_{1,q}|^q)\eta dx \leq 0 \leq \lambda \int_{\bar{\Omega}_\epsilon} c(x)\psi_1^\delta \eta dx + \int_{\bar{\Omega}_\epsilon} d(x)\psi_2^\beta \eta dx.$$

Furthermore, we note that $\phi_{1,p}, \phi_{1,q} \geq \sigma$ in $\Omega - \bar{\Omega}_\epsilon$ for some $\sigma > 0$. Then from (3.2) there is $\xi_0 > 0$ such that if $\xi \in (0, \xi_0)$, the following inequalities hold:

$$\xi^{p-1-k\gamma} \left(\frac{p}{p-1}\right)^{p-1} \lambda_1^{(p)}a(x)\phi_{1,p}^p \leq \lambda a(x) \sigma^{\frac{\gamma q}{q-1}} \leq \lambda a(x) \phi_{1,q}^{\frac{\gamma q}{q-1}} \quad \text{in } \Omega - \bar{\Omega}_\epsilon,$$

and

$$\xi^{k(q-1)-\delta} \left(\frac{q}{q-1}\right)^{q-1} \lambda_1^{(q)}c(x)\phi_{1,q}^q \leq \lambda c(x) \sigma^{\frac{\delta p}{p-1}} \leq \lambda c(x) \phi_{1,p}^{\frac{\alpha p}{p-1}} \quad \text{in } \Omega - \bar{\Omega}_\epsilon.$$

Then, we have

$$\begin{aligned} \int_{\Omega-\bar{\Omega}_\epsilon} P(x)|\nabla\psi_1|^{p-2}\nabla\psi_1\cdot\nabla\zeta dx &= \left(\frac{p}{p-1}\xi\right)^{p-1} \int_{\Omega-\bar{\Omega}_\epsilon} (\lambda_1^{(p)}a(x)\phi_{1,p}^p - P(x)|\nabla\phi_{1,p}|^p)\zeta dx \\ &\leq \lambda \int_{\Omega-\bar{\Omega}_\epsilon} a(x)\xi^{\gamma k}\phi_{1,q}^{\frac{\gamma q}{q-1}}\zeta dx \leq \lambda \int_{\Omega-\bar{\Omega}_\epsilon} a(x)\psi_2^\gamma\zeta dx + \int_{\Omega-\bar{\Omega}_\epsilon} b(x)\psi_1^\alpha\zeta dx. \end{aligned}$$

Similarly,

$$\begin{aligned} \int_{\Omega-\bar{\Omega}_\epsilon} Q(x)|\nabla\psi_2|^{q-2}\nabla\psi_2\cdot\nabla\eta dx &= \left(\frac{q}{q-1}\xi^k\right)^{q-1} \int_{\Omega-\bar{\Omega}_\epsilon} (\lambda_1^{(q)}c(x)\phi_{1,q}^q - Q(x)|\nabla\phi_{1,q}|^q)\eta dx \\ &\leq \lambda \int_{\Omega-\bar{\Omega}_\epsilon} c(x)\xi^\delta\phi_{1,p}^{\frac{\delta p}{p-1}}\eta dx \leq \lambda \int_{\Omega-\bar{\Omega}_\epsilon} c(x)\psi_1^\delta\eta dx + \int_{\Omega-\bar{\Omega}_\epsilon} d(x)\psi_2^\alpha\eta dx, \end{aligned}$$

i.e. (ψ_1, ψ_2) is a subsolution of (1.4).

Next, we construct a supersolution (z_1, z_2) of system (1.4). Let $e_r(x)$ be the positive solution of (see [23])

$$-\Delta_{R,r}e_r = 1 \quad \text{in } \Omega, \quad e_r = 0 \quad \text{on } \partial\Omega \quad \text{for } r = p, q.$$

We denote $z_1(x) = Ae_p$, $z_2(x) = Be_q$, where the constants $A, B > 0$ are large and to be chosen later. We shall verify that (z_1, z_2) is the supersolution of (1.4). To do this, let $\zeta \in W_0^{1,p}(P, \Omega)$ with $\zeta \geq 0$. Then we have

$$\int_{\Omega} P(x)|\nabla z_1|^{p-2}\nabla z_1\cdot\nabla\zeta dx = A^{p-1} \int_{\Omega} P(x)|\nabla e_p|^{p-2}\nabla e_p\cdot\nabla\zeta dx = A^{p-1} \int_{\Omega} \zeta dx.$$

Similarly, for $\eta \in W_0^{1,q}(Q, \Omega)$ with $\eta \geq 0$, we have

$$\int_{\Omega} Q(x)|\nabla z_2|^{q-2}\nabla z_2\cdot\nabla\eta dx = B^{q-1} \int_{\Omega} Q(x)|\nabla e_q|^{q-2}\nabla e_q\cdot\nabla\eta dx = B^{q-1} \int_{\Omega} \eta dx.$$

Since $(p-1)(q-1) - \gamma\delta > 0$, it is easy to prove that there exist positive large constants A, B such that

$$A^{p-1} = \lambda l_a B^\gamma \mu_q^\gamma + l_b A^\alpha \mu_p^\alpha, \quad B^{q-1} = \lambda l_c A^\delta \mu_p^\delta + l_d B^\beta \mu_q^\beta,$$

where $l_a = |a(x)|$, $l_b = |b(x)|$, $l_c = |c(x)|$, $l_d = |d(x)|$ and $\mu_r = \|e_r\|_\infty$; $r = p, q$. These imply that

$$\begin{aligned} \int_{\Omega} P(x)|\nabla z_1|^{p-2}\nabla z_1\cdot\nabla\zeta dx &= \lambda \int_{\Omega} l_a B^\gamma \mu_q^\gamma \zeta dx + \int_{\Omega} l_b A^\alpha \mu_p^\alpha \zeta dx \\ &\geq \lambda \int_{\Omega} a(x)z_2^\gamma \zeta dx + \int_{\Omega} b(x)z_1^\alpha \zeta dx, \end{aligned}$$

and

$$\begin{aligned} \int_{\Omega} Q(x)|\nabla z_2|^{q-2}\nabla z_2\cdot\nabla\eta dx &= \lambda \int_{\Omega} l_c A^\delta \mu_p^\delta \eta dx + \int_{\Omega} l_d B^\beta \mu_q^\beta \eta dx \\ &\geq \lambda \int_{\Omega} c(x)z_1^\delta \eta dx + \int_{\Omega} d(x)z_2^\beta \eta dx, \end{aligned}$$

i.e. (z_1, z_2) is a supersolution of (1.4) with $z_i \geq \psi_i$ with large A, B , for $i = 1, 2$. Thus, there exists a solution (u, v) of (1.4) with $\psi_1 \leq u \leq z_1$, $\psi_2 \leq v \leq z_2$. This completes the proof. \square

4. NONEXISTENCE RESULT

In this section, under some certain conditions we prove that system (1.4) has no positive weak solution.

Theorem 4.1. *Suppose that $(p - 1)(q - 1) - \gamma\delta = 0$, $p\gamma = q(p - 1)$ and $b(x), d(x) \leq a(x) = c(x)$ for all $x \in \Omega$. Then there exists $\lambda_0 > 0$ such that for $0 < \lambda < \lambda_0$, system (1.4) has no positive weak solution.*

Proof. Let us assume that $(u, v) \in W_0^{1,p}(P, \Omega) \times W_0^{1,p}(Q, \Omega)$ be a positive weak solution of (1.4). We prove this theorem by arriving at a contradiction.

Multiplying the first equation of (1.4) by u , we have from Young inequality that

$$(4.1) \quad \int_{\Omega} P(x)|\nabla u|^p dx \leq \lambda \int_{\Omega} a(x) \left(\frac{u^p}{\mu_1} + \frac{v^q}{\mu_2} \right) dx + \int_{\Omega} b(x)u^p,$$

with $\mu_1 = p > 1$ and $\mu_2 = \frac{p}{p-1} > 1$.

Similarly, we have

$$(4.2) \quad \int_{\Omega} Q(x)|\nabla v|^q dx \leq \lambda \int_{\Omega} c(x) \left(\frac{u^p}{\theta_1} + \frac{v^q}{\theta_2} \right) dx + \int_{\Omega} d(x)v^q,$$

with $\theta_1 = \frac{q}{q-1} > 1$ and $\theta_2 = q > 1$.

Note that

$$(4.3) \quad \lambda_1^{(p)} \int_{\Omega} a(x)u^p dx \leq \int_{\Omega} P(x)|\nabla u|^p dx, \quad \lambda_1^{(q)} \int_{\Omega} c(x)v^q dx \leq \int_{\Omega} Q(x)|\nabla v|^q dx.$$

Combining (4.1)-(4.3), we obtain

$$\lambda_1^{(p)} \int_{\Omega} a(x)u^p dx + \lambda_1^{(q)} \int_{\Omega} c(x)v^q dx \leq \lambda \int_{\Omega} \left(\frac{a(x)}{\mu_1} + \frac{c(x)}{\theta_1} \right) u^p dx + \lambda \int_{\Omega} \left(\frac{a(x)}{\mu_2} + \frac{c(x)}{\theta_2} \right) v^q dx + \int_{\Omega} b(x)u^p + \int_{\Omega} d(x)v^q.$$

Now, if $b(x), d(x) \leq a(x) = c(x)$ for all $x \in \Omega$, we have

$$(\lambda_1^{(p)} - 1 - \lambda) \int_{\Omega} a(x)u^p dx + (\lambda_1^{(q)} - 1 - \lambda) \int_{\Omega} a(x)v^q dx \leq 0,$$

which is a contradiction if $0 < \lambda < \lambda_0 = \min\{\lambda_1^{(p)} - 1, \lambda_1^{(q)} - 1\}$. This completes the proof. \square

Acknowledgement. The author would like to express his gratitude to Professor H. M. Serag (Mathematics Department, Faculty of Science, AL- Azhar University) for continuous encouragement during the development of this work.

REFERENCES

- [1] J. Ali and R. Shivaji, *Positive solutions for a class of p -Laplacian systems with multiple parameters*, J. Math. Anal. Appl., **335** (2007), 1013-1019.
- [2] M. Alimohmmady and M. Koozegar, *Some remarks on the nonexistence of positive solutions for some a, p -Laplacian systems*, The Journal of Nonlinear Sciences and Applications, **1** (2008), 56-60.
- [3] A. Canada, P. Dravek and J. Gamez, *Existence of positive solutions for some problems with nonlinear diffusion*, Trans. Amer. Math. Soc., **349** (1997), 4231-4249.
- [4] F. Cirstea and D. Radulescu, *Existence and Non-existence Results for a Quasilinear Problem with Nonlinear Boundary Condition*, Journal of Mathematical Analysis and Applications, **244** (2000,) 169-183.
- [5] F. Cirstea, et al, *Weak solutions of quasilinear problems with nonlinear boundary condition*, Nonlinear Analysis, **43** (2001), 623-636.

- [6] P. Drabek, A. Kufner and F. Nicolosi, *Quasilinear elliptic equation with degenerations and singularities*, Walter de Gruyter, Bertin, New York, 1997.
- [7] S. Khafagy, *Existence and nonexistence of positive weak solutions for a class of (p, q) -Laplacian nonlinear elliptic system with different weights*, Int. J. Contemp. Math. Sciences, **6** (2011), 2391-2400.
- [8] S. Khafagy, *Existence, nonexistence and uniqueness of positive weak solution for a nonlinear system involving weighted p -Laplacian*, Global Journal of Pure and Applied Mathematics, **8** (2012), 205-214.
- [9] S. Khafagy, *Existence and nonexistence of positive weak solutions for a class of weighted (p, q) -Laplacian nonlinear system*, Global Journal of Pure and Applied Mathematics, **9** (2013), 379-387.
- [10] S. Khafagy, *Maximum Principle and Existence of Weak Solutions for Nonlinear System Involving Weighted (p, q) -Laplacian*, Southeast Asian Bulletin of Mathematics, **40** (2016), 353-364.
- [11] S. Khafagy, *Maximum Principle and Existence of Weak Solutions for Nonlinear System Involving Singular p -Laplacian Operators*, Journal of Partial differential equations, **29** (2016), 89-101.
- [12] S. Khafagy, *Non-existence of positive weak solutions for some weighted p -Laplacian system*, Journal of Advanced Research in Dynamical and Control Systems, **7** (2015), 71-77.
- [13] S. Khafagy, *On the stability of positive weak solution for weighted p -Laplacian nonlinear system*, New Zealand Journal of Mathematics, **45** (2015), 39-43.
- [14] S. Khafagy, *On positive weak solutions for nonlinear elliptic system involving singular p -Laplacian operator*, Journal of Mathematical Analysis, **7** (2016), 10-17.
- [15] S. Khafagy, *On positive weak solution for a nonlinear system involving weighted p -Laplacian*, Journal of Advanced Research in Dynamical and Control Systems, **4** (2012), 50-58.
- [16] S. Khafagy and H. Serag, *Existence of Weak Solutions for $n \times n$ Nonlinear Systems Involving Different p -Laplacian Operators*, Electron. J. Diff. Eqns., **2009** (2009), 1-14.
- [17] A. Leung, *Systems of nonlinear partial differential equations. Applications to biology and engineering*, Math. Appl. (Kluwer Academic Publishers, Dordrecht, 1989).
- [18] A. Leung, *Nonlinear Systems of Partial Differential Equations: Applications to Life and Physical Sciences*, World Scientific Publishing Co., 2009
- [19] C. Pao, *Nonlinear Parabolic and Elliptic Equations*, Plenum Press, New York, 1992.
- [20] K. Pflueger, *Existence and multiplicity of solutions to a p -Laplacian equation with nonlinear boundary condition*, Electron. J. Diff. Eqns., **1998** (1998), 1-13.
- [21] H. Serag and S. Khafagy, *Existence of Weak Solutions for $n \times n$ Nonlinear Systems Involving Different Degenerated p -Laplacian Operators*, New Zealand Journal of Mathematics, **38** (2008), 75-86.
- [22] H. Serag and S. Khafagy, *On a nonhomogeneous elliptic systems $n \times n$ Involving p -Laplacian with different weights*, Journal of Advanced Research in Differential Equations, **1** (2009), 47-62.
- [23] H. Serag and S. Khafagy, *On Maximum Principle and Existence of Positive Weak Solutions for $n \times n$ Nonlinear Systems Involving Degenerated p -Laplacian Operator*, Turk J M, **34** (2010), 59-71.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, AL-AZHAR UNIVERSITY, NASR CITY (11884), CAIRO, EGYPT.

CURRENT ADDRESS: DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE IN ZULFI, MAJMAAH UNIVERSITY, ZULFI 11932, P.O. BOX 1712, SAUDI ARABIA

E-mail address: el.gharieb@hotmail.com, s.khafagy@mu.edu.sa