

DEPTH AND STANLEY DEPTH OF THE EDGE IDEALS OF SOME m -LINE GRAPHS AND m -CYCLIC GRAPHS WITH A COMMON VERTEX

GUANGJUN ZHU

ABSTRACT. We give some precise formulas for the depth of the quotient rings of the edge ideals associated to a graph consisting, either of the union of some line graphs $L_{3r_1}, \dots, L_{3r_{k_1}}, L_{3s_1+1}, \dots, L_{3s_{k_2}+1}$ and $L_{3t_1+2}, \dots, L_{3t_{k_3}+2}$ or of the union of cycle graphs $C_{3r_1}, \dots, C_{3r_{k_1}}, C_{3s_1+1}, \dots, C_{3s_{k_2}+1}$ and $C_{3t_1+2}, \dots, C_{3t_{k_3}+2}$, with a common vertex. We also give some tight bounds for their Stanley depths.

Mathematics Subject Classification (2010): 13C15; 13P10; 13F20.

Keywords: Stanley depth, depth, Stanley conjecture, edge ideal.

Article history:

Received 18 October 2015

Received in revised form 4 November 2015

Accepted 6 November 2015

1. INTRODUCTION

Let $S = K[x_1, \dots, x_n]$ be the polynomial ring in n variables over a field K and M a finitely generated \mathbb{Z}^n -graded S -module. For a homogeneous element $u \in M$ and a subset $Z \subseteq \{x_1, \dots, x_n\}$, $uK[Z]$ denotes the K -subspace of M generated by all the homogeneous elements of the form uv , where v is a monomial in $K[Z]$. The \mathbb{Z}^n -graded K -subspace $uK[Z]$ is said to be a Stanley space of dimension $|Z|$ if it is a free $K[Z]$ -module, where, as usual, $|Z|$ denotes the number of elements of Z . A Stanley decomposition of M is a decomposition of M as a finite direct sum of \mathbb{Z}^n -graded K -vector spaces

$$\mathcal{D} : M = \bigoplus_{i=1}^r u_i K[Z_i]$$

where each $u_i K[Z_i]$ is a Stanley space of M . The number

$$\text{sdepth}(\mathcal{D}) = \min\{|Z_i| : i = 1, \dots, r\}$$

is called the Stanley depth of decomposition \mathcal{D} and the quantity

$$\text{sdepth}(M) := \max\{\text{sdepth}(\mathcal{D}) \mid \mathcal{D} \text{ is a Stanley decomposition of } M\}$$

We greatly acknowledge the use of the computer algebra system CoCoA ([3]) for our experiments.

Supported by the National Natural Science Foundation of China (11271275) and by Foundation of Jiangsu Overseas Research & Training Program for University Prominent Young & Middle-aged Teachers and Presidents and by Foundation of the Priority Academic Program Development of Jiangsu Higher Education Institutions.

is called the Stanley depth of M . Stanley [10] conjectured that

$$\text{sdepth}(M) \geq \text{depth}(M)$$

for all \mathbb{Z}^n -graded S -modules M . This conjecture proves to be false, in general, for $M = S/I$ and $M = J/I$, where $I \subset J \subset S$ are monomial ideals, see [4].

Herzog, Vlădoiu and Zheng [6] introduced a method to compute the Stanley depth of a factor of two monomial ideals which was later developed into an effective algorithm by Rinaldo [9] implemented in CoCoA [3]. However, it is difficult to compute this invariant, even in some very particular cases. For instance in [1] Biró et al. proved that $\text{sdepth}(\mathfrak{m}) = \lceil \frac{n}{2} \rceil$ where $\mathfrak{m} = (x_1, \dots, x_n)$ is the graded maximal ideal of S and $\lceil \frac{n}{2} \rceil$ denote the smallest integer $\geq \frac{n}{2}$. For a friendly introduction on Stanley depth we refer the reader to [5].

Let I_n and J_n be the edge ideals associated to the line, respectively, cycle graph of length n . Morey [7] proved that $\text{depth}(S/I_n) = \lceil \frac{n}{3} \rceil$. Replacing depth by stanley depth, Ştefan [11] showed that $\text{sdepth}(S/I_n) = \lceil \frac{n}{3} \rceil$. In [2], Cimpoeaş proved that $\text{depth}(S/J_n) = \lceil \frac{n-1}{3} \rceil$ and $\text{sdepth}(S/J_n) = \lceil \frac{n-1}{3} \rceil$ for $n \equiv 0 \pmod{3}$ and $n \equiv 2 \pmod{3}$. He also proved that $\lceil \frac{n-1}{3} \rceil \leq \text{sdepth}(S/J_n) \leq \lceil \frac{n}{3} \rceil$ for $n \equiv 1 \pmod{3}$. Let I and J be the edge ideals associated to the graph consisting of the union of line graphs $L_{3r_1}, \dots, L_{3r_{k_1}}, L_{3s_1+1}, \dots, L_{3s_{k_2}+1}$ and $L_{3t_1+2}, \dots, L_{3t_{k_3}+2}$ with a common vertex, respectively, the graph consisting of the union of cycle graphs $C_{3r_1}, \dots, C_{3r_{k_1}}, C_{3s_1+1}, \dots, C_{3s_{k_2}+1}$ and $C_{3t_1+2}, \dots, C_{3t_{k_3}+2}$ with a common vertex, then using similar techniques, we prove that

$$(1) \text{ sdepth}\left(\frac{S}{I}\right) \geq \text{depth}\left(\frac{S}{I}\right) = \begin{cases} \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i, & \text{if } k_3 = 0, \\ \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + 1, & \text{otherwise;} \end{cases}$$

$$(2) \text{ sdepth}\left(\frac{S}{I}\right) \leq \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + 1.$$

In the fourth section, we prove that

$$(1) \text{ sdepth}\left(\frac{S}{J}\right) \geq \text{depth}\left(\frac{S}{J}\right) = \begin{cases} \sum_{i=1}^{k_2} s_i, & \text{if } k_1 = k_3 = 0, \\ \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1, & \text{otherwise;} \end{cases}$$

$$(2) \text{ sdepth}\left(\frac{S}{J}\right) \leq \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1;$$

$$(3) \text{ sdepth}\left(\frac{J}{I}\right) \geq \text{depth}\left(\frac{J}{I}\right) = \begin{cases} \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_3} t_i - k_1 + 2, & \text{if } k_2 = 0, \\ \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1, & \text{otherwise.} \end{cases}$$

2. PRELIMINARIES

We first recall some definitions about graphs and their edge ideals in order to make this paper self-contained. However, for more details on the notions, we refer the reader to [13, 14].

Definition 2.1. Let $G_i = (V(G_i), E(G_i))$ be some graphs with vertex set $V(G_i)$ and edge set $E(G_i)$ for $1 \leq i \leq k$. The union of graphs G_1, G_2, \dots, G_k , written as $G_1 \cup G_2 \cup \dots \cup G_k$, is the graph with vertex set $\bigcup_{i=1}^k V(G_i)$ and edge set $\bigcup_{i=1}^k E(G_i)$.

Definition 2.2. Let $G = (V(G), E(G))$ be a graph with vertex set $V(G) = \{x_1, \dots, x_n\}$ and edge set $E(G)$. Suppose that x_1, \dots, x_n are variables over the field K . The edge ideal of graph G in the polynomial ring $S = K[x_1, \dots, x_n]$ is the squarefree monomial ideal

$$I(G) = (\{x_i x_j \mid \{x_i, x_j\} \in E(G)\}).$$

For the sake of simplicity, we will use the same notation $x_i x_j$ for the monomial and for the corresponding edge of graph G .

Definition 2.3. Let $G = (V(G), E(G))$ be a graph with vertex set $V(G) = \{x_1, \dots, x_m\}$ and edge set $E(G)$. Then G is called a line graph of length m , denoted by L_m , if its edge set $E = \{x_i x_{i+1} \mid 1 \leq i \leq m-1\}$. Similarly, if $m \geq 3$, then G is called a cycle graph of length m , denoted by C_m , if its edge set $E = \{x_i x_{i+1} \mid 1 \leq i \leq m-1\} \cup \{x_m x_1\}$.

We recall the well known Depth Lemma, see for instance [13, Lemma 1.3.9] or [12, Lemma 3.1.4].

Lemma 2.4. (*Depth Lemma*) Let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be a short exact sequence of modules over a local ring S , or a Noetherian graded ring with S_0 local, then

- (i) $\text{depth}(M) \geq \min\{\text{depth}(L), \text{depth}(N)\}$;
- (ii) $\text{depth}(L) \geq \min\{\text{depth}(M), \text{depth}(N) + 1\}$;
- (iii) $\text{depth}(N) \geq \min\{\text{depth}(L) - 1, \text{depth}(M)\}$.

The most of the statements of the Depth Lemma are wrong if we replace depth by Stanley depth. Rauf [8] proved the analog of Lemma 2.4 (i) for Stanley depth.

Lemma 2.5. Let $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ be a short exact sequence of finitely generated \mathbb{Z}^n -graded S -modules. Then

$$s\text{depth}(M) \geq \min\{s\text{depth}(L), s\text{depth}(N)\}.$$

Using Depth Lemma, Morey in [7] proved the following result.

Lemma 2.6. Let L_m be a line graph of length m and $I(L_m)$ its edge ideal. Then $\text{depth}(S/I(L_m)) = \lceil \frac{m}{3} \rceil$.

Replacing depth by Stanley depth, Ştefan in [11] showed the following result.

Lemma 2.7. Let L_m be a line graph of length m and $I(L_m)$ its edge ideal. Then $s\text{depth}(S/I(L_m)) = \lceil \frac{m}{3} \rceil$.

3. DEPTH AND STANLEY DEPTH OF THE EDGE IDEALS OF SOME m -LINE GRAPHS WITH A COMMON VERTEX

In this section, we will give some formulas for depth and Stanley depth of the quotient rings of the edge ideals of some m -line graphs with a common vertex. We assume that G is the m -line graph formed by joining m line graphs $L_{3r_1}, \dots, L_{3r_{k_1}}, L_{3s_1+1}, \dots, L_{3s_{k_2}+1}, L_{3t_1+2}, \dots, L_{3t_{k_3}+2}$ at a common vertex, where $k_1 + k_2 + k_3 = m$ and $k_i \geq 0$ for $i = 1, 2, 3$. We adopt the following notation to edges of graph G :

$$E(L_{3r_i, i}) = \{x_1 x_{2, i}, x_{2, i} x_{3, i}, \dots, x_{3r_i-1, i} x_{3r_i, i}\} \text{ for any } 1 \leq i \leq k_1 \text{ and } r_1 \leq \dots \leq r_{k_1},$$

$$E(L_{3s_i+1, i}) = \{x_1 y_{2, i}, y_{2, i} y_{3, i}, \dots, y_{3s_i, i} y_{3s_i+1, i}\} \text{ for any } 1 \leq i \leq k_2 \text{ and } s_1 \leq \dots \leq s_{k_2},$$

$$E(L_{3t_i+2, i}) = \{x_1 z_{2, i}, z_{2, i} z_{3, i}, \dots, z_{3t_i+1, i} z_{3t_i+2, i}\} \text{ for all } 1 \leq i \leq k_3 \text{ and } t_1 \leq \dots \leq t_{k_3}.$$

Set K be any field, $S = K[x_1, x_{2,1}, \dots, x_{3r_1,1}, \dots, x_{2,k_1}, \dots, x_{3r_{k_1},k_1}, y_{2,1}, \dots, y_{3s_1+1,1}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}, z_{2,1}, \dots, z_{3t_1+2,1}, \dots, z_{2,k_3}, \dots, z_{3t_{k_3}+2,k_3}]$ the polynomial ring. The edge ideal of graph G is $I = (x_1 x_{2,1}, x_{2,1} x_{3,1}, \dots, x_{3r_1-1,1} x_{3r_1,1}, \dots, x_1 x_{2,k_1}, x_{2,k_1} x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1} x_{3r_{k_1},k_1}, x_1 y_{2,1}, y_{2,1} y_{3,1}, \dots, y_{3s_1,1} y_{3s_1+1,1}, \dots, x_1 y_{2,k_2}, y_{2,k_2} y_{3,k_2}, \dots, y_{3s_{k_2},k_2} y_{3s_{k_2}+1,k_2}, x_1 z_{2,1}, z_{2,1} z_{3,1}, \dots, z_{3t_1+1,1} z_{3t_1+2,1}, \dots, x_1 z_{2,k_3}, \dots, z_{3t_{k_3}+1,k_3} z_{3t_{k_3}+2,k_3})$.

Example 3.1. The following graph G is the union of 5 line graphs L_3, L_4, L_5, L_6 and L_7 with a common vertex x_1 .




Figure 1

The edge ideal of graph G is $I = (x_1x_{2,1}, x_{2,1}x_{3,1}, x_1x_{2,2}, x_{2,2}x_{3,2}, x_{3,2}x_{4,2}, x_{4,2}x_{5,2}, x_{5,2}x_{6,2}, x_1y_{2,1}, y_{2,1}y_{3,1}, y_{3,1}y_{4,1}, x_1y_{2,2}, y_{2,2}y_{3,2}, y_{3,2}y_{4,2}, y_{4,2}y_{5,2}, y_{5,2}y_{6,2}, y_{6,2}y_{7,2}, x_1z_{2,1}, z_{2,1}z_{3,1}, z_{3,1}z_{4,1}, z_{4,1}z_{5,1})$.

We need the following lemma (See [8, Theorem 3.1]).

Lemma 3.2. Let $I \subset S_1 = K[x_1, \dots, x_m]$, $J \subset S_2 = K[y_1, \dots, y_n]$ be monomial ideals and $S = K[x_1, \dots, x_m, y_1, \dots, y_n]$. Then

$$sdepth(S/(IS, JS)) \geq sdepth(S_1/IS_1) + sdepth(S_2/JS_2).$$

Now, we prove the main results of this section. We adopt the following convention: whenever, in a sum, the index runs from 1 to 0, the sum has to be taken equal to zero.

Theorem 3.3. Let G be a graph consisting of the union of line graphs $L_{3r_1}, \dots, L_{3r_{k_1}}, L_{3s_1+1}, \dots, L_{3s_{k_2}+1}$ and $L_{3t_1+2}, \dots, L_{3t_{k_3}+2}$ with a common vertex x_1 , where $k_i \geq 0$ for $i = 1, 2, 3$. Let I be its edge ideal. Then

$$sdepth\left(\frac{S}{I}\right) \geq depth\left(\frac{S}{I}\right) = \begin{cases} \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i, & \text{if } k_3 = 0, \\ \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + 1, & \text{otherwise.} \end{cases}$$

In particular, S/I satisfies the Stanley conjecture.

Proof. Note that $(I : x_1) = (x_{2,1}, \dots, x_{2,k_1}, y_{2,1}, \dots, y_{2,k_2}, z_{2,1}, \dots, z_{2,k_3}, x_{3,1}x_{4,1}, \dots, x_{3r_1-1,1}x_{3r_1,1}, \dots, x_{3,k_1}x_{4,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1}, y_{3,1}y_{4,1}, \dots, y_{3s_1,1}y_{3s_1+1,1}, \dots, y_{3,k_2}y_{4,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2}, z_{3,1}z_{4,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1}, \dots, z_{3,k_3}z_{4,k_3}, \dots, z_{3t_{k_3}+1,k_3}z_{3t_{k_3}+2,k_3})$ and $(I, x_1) = (x_{2,1}x_{3,1}, \dots, x_{3r_1-1,1}x_{3r_1,1}, \dots, x_{2,k_1}x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1}, y_{2,1}y_{3,1}, \dots, y_{3s_1,1}y_{3s_1+1,1}, \dots, y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2}, z_{2,1}z_{3,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1}, \dots, z_{2,k_3}z_{3,k_3}, \dots, z_{3t_{k_3}+1,k_3}z_{3t_{k_3}+2,k_3}, x_1)$, thus we get that

$$\begin{aligned} \frac{S}{(I : x_1)} &\cong \frac{K[x_{3,1}, \dots, x_{3r_1,1}]}{(x_{3,1}x_{4,1}, \dots, x_{3r_1-1,1}x_{3r_1,1})} \otimes_K \cdots \otimes_K \frac{K[x_{3,k_1}, \dots, x_{3r_{k_1},k_1}]}{(x_{3,k_1}x_{4,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1})} \\ &\otimes_K \frac{K[y_{3,1}, \dots, y_{3s_1+1,1}]}{(y_{3,1}y_{4,1}, \dots, y_{3s_1,1}y_{3s_1+1,1})} \otimes_K \cdots \otimes_K \frac{K[y_{3,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{3,k_2}y_{4,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})} \\ &\otimes_K \frac{K[z_{3,1}, \dots, z_{3t_1+2,1}]}{(z_{3,1}z_{4,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1})} \otimes_K \cdots \otimes_K \frac{K[z_{3,k_3}, \dots, z_{3t_{k_3}+2,k_3}]}{(z_{3,k_3}z_{4,k_3}, \dots, z_{3t_{k_3}+1,k_3}z_{3t_{k_3}+2,k_3})} \otimes_K K[x_1], \end{aligned}$$

and

$$\begin{aligned} \frac{S}{(I, x_1)} &\cong \frac{K[x_{2,1}, \dots, x_{3r_1,1}]}{(x_{2,1}x_{3,1}, \dots, x_{3r_1-1,1}x_{3r_1,1})} \otimes_K \cdots \otimes_K \frac{K[x_{2,k_1}, \dots, x_{3r_{k_1},k_1}]}{(x_{2,k_1}x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1})} \\ &\otimes_K \frac{K[y_{2,1}, \dots, y_{3s_1+1,1}]}{(y_{2,1}y_{3,1}, \dots, y_{3s_1,1}y_{3s_1+1,1})} \otimes_K \cdots \otimes_K \frac{K[y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})} \\ &\otimes_K \frac{K[z_{2,1}, \dots, z_{3t_1+2,1}]}{(z_{2,1}z_{3,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1})} \otimes_K \cdots \otimes_K \frac{K[z_{2,k_3}, \dots, z_{3t_{k_3}+2,k_3}]}{(z_{2,k_3}z_{3,k_3}, \dots, z_{3t_{k_3}+1,k_3}z_{3t_{k_3}+2,k_3})}. \end{aligned}$$

Therefore, by Lemmas 2.6, 2.7, 3.2 and [13, Proposition 2.2.20, Theorem 2.2.21], we obtain that $\text{sdepth}(\frac{S}{(I, x_1)}) \geq \text{depth}(\frac{S}{(I, x_1)}) = \sum_{i=1}^{k_1} \lceil \frac{3r_i-2}{3} \rceil + \sum_{i=1}^{k_2} \lceil \frac{3s_i-1}{3} \rceil + \sum_{i=1}^{k_3} \lceil \frac{3t_i}{3} \rceil + 1 = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + 1$, and $\text{sdepth}(\frac{S}{(I, x_1)}) \geq \text{depth}(\frac{S}{(I, x_1)}) = \sum_{i=1}^{k_1} \lceil \frac{3r_i-1}{3} \rceil + \sum_{i=1}^{k_2} \lceil \frac{3s_i}{3} \rceil + \sum_{i=1}^{k_3} \lceil \frac{3t_i+1}{3} \rceil = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + k_3$.

Using Lemma 2.5 on the short exact sequence

$$(1) \quad 0 \longrightarrow S/(I : x_1) \xrightarrow{x_1} S/I \longrightarrow S/(I, x_1) \longrightarrow 0,$$

we conclude that $\text{sdepth}(\frac{S}{I}) \geq \begin{cases} \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i, & \text{if } k_3 = 0; \\ \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + 1, & \text{otherwise.} \end{cases}$

If $k_3 \neq 0$, then $\text{depth}(\frac{S}{(I, x_1)}) \geq \text{depth}(\frac{S}{(I, x_1)})$. Using Lemma 2.4 on the short exact sequence (1), it follows that $\text{depth}(\frac{S}{(I, x_1)}) = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + 1$.

Assume that $k_3 = 0$. We claim that we have the S -module isomorphism

$$\begin{aligned} \frac{(I : x_1)}{I} &\cong \bigoplus_{i=1}^{k_1} x_{2,i} \left(\frac{K[x_{3,1}, \dots, x_{3r_1,1}]}{(x_{3,1}x_{4,1}, \dots, x_{3r_1-1,1}x_{3r_1,1})} \otimes_K \cdots \otimes_K \frac{K[x_{3,i-1}, \dots, x_{3r_{i-1},i-1}]}{(x_{3,i-1}x_{4,i-1}, \dots, x_{3r_{i-1}-1,i-1}x_{3r_{i-1},i-1})} \right. \\ &\otimes_K \frac{K[x_{4,i}, \dots, x_{3r_i,i}]}{(x_{4,i}x_{5,i}, \dots, x_{3r_i-1,i}x_{3r_i,i})} \otimes_K \frac{K[x_{2,i+1}, \dots, x_{3r_{i+1},i+1}]}{(x_{2,i+1}x_{3,i+1}, \dots, x_{3r_{i+1}-1,i+1}x_{3r_{i+1},i+1})} \otimes_K \cdots \\ &\otimes_K \frac{K[x_{2,k_1}, \dots, x_{3r_{k_1},k_1}]}{(x_{2,k_1}x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1})} \otimes_K \frac{K[y_{2,1}, \dots, y_{3s_1+1,1}]}{(y_{2,1}y_{3,1}, \dots, y_{3s_1,1}y_{3s_1+1,1})} \otimes_K \cdots \\ &\otimes_K \frac{K[y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})} \Big) [x_{2,i}] \oplus \left(\bigoplus_{i=1}^{k_2} y_{2,i} \left(\frac{K[x_{3,1}, \dots, x_{3r_1,1}]}{(x_{3,1}x_{4,1}, \dots, x_{3r_1-1,1}x_{3r_1,1})} \otimes_K \cdots \right. \right. \\ &\otimes_K \frac{K[x_{3,k_1}, \dots, x_{3r_{k_1},k_1}]}{(x_{3,k_1}x_{4,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1})} \otimes_K \frac{K[y_{3,1}, \dots, y_{3s_1+1,1}]}{(y_{3,1}y_{4,1}, \dots, y_{3s_1,1}y_{3s_1+1,1})} \otimes_K \cdots \\ &\otimes_K \frac{K[y_{3,i-1}, \dots, y_{3s_{i-1}+1,i-1}]}{(y_{3,i-1}y_{4,i-1}, \dots, y_{3s_{i-1}-1,i-1}y_{3s_{i-1}+1,i-1})} \otimes_K \frac{K[y_{4,i}, \dots, y_{3s_i+1,i}]}{(y_{4,i}y_{5,i}, \dots, y_{3s_i,i}y_{3s_i+1,i})} \otimes_K \cdots \\ &\left. \left. \otimes_K \frac{K[y_{2,i+1}, \dots, y_{3s_{i+1}+1,i+1}]}{(y_{2,i+1}y_{3,i+1}, \dots, y_{3s_{i+1},i+1}y_{3s_{i+1}+1,i+1})} \otimes_K \cdots \otimes_K \frac{K[y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})} \right) [y_{2,i}], \end{aligned}$$

where $x_{i,0} = y_{j,0} = 0$ for $3 \leq i \leq k_1$, $3 \leq j \leq k_2$. Indeed, if $u \in (I : x_1)$ is a monomial such that $u \notin I$, then $x_{2,i}|u$ or $y_{2,j}|u$ for some $1 \leq i \leq k_1$ or $1 \leq j \leq k_2$.

If $x_{2,1}|u$, then we can write u as $u = x_{2,1}^\alpha v$ with $\alpha \geq 1$ and $x_{2,1} \nmid v$. Since $u \notin I$, we have that $v \in K[x_{4,1}, \dots, x_{3r_1,1}, x_{2,2}, \dots, x_{3r_2,2}, \dots, x_{2,k_1}, \dots, x_{3r_{k_1},k_1}, y_{2,1}, \dots, y_{3s_1+1,1}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]$ and

$v \notin (x_{4,1}x_{5,1}, \dots, x_{3r_1-1,1}x_{3r_1,1}, x_{2,2}x_{3,2}, \dots, x_{3r_2-1,2}x_{3r_2,2}, \dots, x_{2,k_1}x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1}, y_{2,1}y_{3,1}, \dots, y_{3s_1,1}y_{3s_1+1,1}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})$. Similarly, if $x_{2,2}|u$ and $x_{2,1} \nmid u$, then $u = x_{2,2}^\alpha v$ with $\alpha \geq 1$ and $v \in K[x_{3,1}, \dots, x_{3r_1,1}, x_{4,2}, \dots, x_{3r_2,2}, x_{2,3}, \dots, x_{3r_3,3}, \dots, x_{2,k_1}, \dots, x_{3r_{k_1},k_1}, y_{2,1}, \dots, y_{3s_1+1,1}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]$ and $v \notin (x_{3,1}x_{4,1}, \dots, x_{3r_1-1,1}x_{3r_1,1}, x_{4,2}x_{5,2}, \dots, x_{3r_2-1,2}x_{3r_2,2}, x_{2,3}x_{3,3}, \dots, x_{3r_3-1,3}x_{3r_3,3}, \dots, x_{2,k_1}x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1}, y_{2,1}y_{3,1}, \dots, y_{3s_1,1}y_{3s_1+1,1}, \dots, y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})$. Other cases can be shown in a similar way as the above.

Therefore, by [13, Proposition 2.2.20, Theorem 2.2.21] and Lemma 2.6, it follows that $\text{depth}(\frac{I:x_1}{I}) = \sum_{i=1}^{k_1-1} \lceil \frac{3r_i-2}{3} \rceil + \lceil \frac{3r_{k_1}-3}{3} \rceil + \sum_{i=1}^{k_2-1} \lceil \frac{3s_i-1}{3} \rceil + \lceil \frac{3s_{k_2}-2}{3} \rceil + 1 = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i$.

Now, using Lemma 2.4 on the short exact sequence

$$(2) \quad 0 \longrightarrow (I : x_1)/I \longrightarrow S/I \longrightarrow S/(I : x_1) \longrightarrow 0,$$

this completes the proof. \square

Assume $n = \sum_{i=1}^{k_1} 3r_i + \sum_{i=1}^{k_2} (3s_i + 1) + \sum_{i=1}^{k_3} (3t_i + 2) - (k_1 + k_2 + k_3) + 1$. We identify S/I with the \mathbb{Z}^n -graded K -subvector space I^c of S which is generated by all monomials $u \in S \setminus I$. Set the set

$$P = \{a \in \mathbb{N}^n : x^a \in I^c \text{ and } x^a | x_1 \prod_{\substack{2 \leq i \leq 3r_j, \\ 1 \leq j \leq k_1}} x_{i,j} \prod_{\substack{2 \leq i \leq 3s_j + 1, \\ 1 \leq j \leq k_2}} y_{i,j} \prod_{\substack{1, 2 \leq i \leq 3t_j + 2, \\ 1 \leq j \leq k_3}} z_{i,j}\},$$

where $a = (a(1), a(2, 1), \dots, a(3r_1, 1), \dots, a(2, k_1), \dots, a(3r_{k_1}, k_1), b(2, 1), \dots, b(3s_1 + 1, 1), \dots, b(2, k_2), \dots, b(3s_{k_2} + 1, k_2), c(2, 1), \dots, c(3t_1 + 2, 1), \dots, c(2, k_3), \dots, c(3t_{k_3} + 2, k_3)) \in \mathbb{N}^n$ and

$$x^a = x_1^{a(1)} \prod_{\substack{2 \leq i \leq 3r_j, \\ 1 \leq j \leq k_1}} x_{i,j}^{a(i,j)} \prod_{\substack{2 \leq i \leq 3s_j + 1, \\ 1 \leq j \leq k_2}} y_{i,j}^{b(i,j)} \prod_{\substack{1, 2 \leq i \leq 3t_j + 2, \\ 1 \leq j \leq k_3}} z_{i,j}^{c(i,j)}.$$

Consider the natural partial order on \mathbb{N}^n which is given by componentwise comparison, i.e. if $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$, then $a \geq b$ if and only if $a_i \geq b_i$ for all $i = 1, \dots, n$. With respect to the partial order induced on P , it becomes a poset where $a \geq a'$ if and only if $x^{a'} | x^a$.

Let $\mathcal{P} : P = \bigcup_{i=1}^r [F_i, G_i]$ be a partition of P . We denote $\text{sdepth}(\mathcal{P}) = \min\{|G_i| : 1 \leq i \leq r\}$. Also, we define the Stanley depth of P , to be the number

$$\text{sdepth}(P) = \max\{\text{sdepth}(\mathcal{P}) \mid \mathcal{P} \text{ is a partition of } P\}.$$

Herzog, Vlădoiu and Zheng proved in [6] that $\text{sdepth}(\frac{S}{I}) = \text{sdepth}(P)$. Now, for $d \in \mathbb{N}$ and $\sigma \in P$, we denote

$$\mathcal{P}_d := \{a \in P : |a| = d\} \quad \text{and} \quad \mathcal{P}_{d,\sigma} := \{a \in \mathcal{P}_d : x^\sigma | x^a\},$$

where for $a = (a(1), a(2, 1), \dots, a(3r_1, 1), \dots, a(2, k_1), \dots, a(3r_{k_1}, k_1), b(2, 1), \dots, b(3s_1 + 1, 1), \dots, b(2, k_2), \dots, b(3s_{k_2} + 1, k_2), c(2, 1), \dots, c(3t_1 + 2, 1), \dots, c(2, k_3), \dots, c(3t_{k_3} + 2, k_3)) \in \mathbb{N}^n$, $|a| := a(1) + \sum_{j=1}^{k_1} \sum_{i=2}^{3r_j} a(i, j) +$

$$\sum_{j=1}^{k_2} \sum_{i=2}^{3s_j+1} b(i, j) + \sum_{j=1}^{k_3} \sum_{i=2}^{3t_j+2} c(i, j).$$

With these notations, we are able to prove the following result.

Theorem 3.4. *Let G be a graph as in Theorem 3.3 and I be its edge ideal. Then*

$$sdepth\left(\frac{S}{I}\right) \leq \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + 1.$$

Proof. Firstly, we claim: if $\sigma \in \mathcal{P}$ such that $\mathcal{P}_{d,\sigma} = \emptyset$, then $sdepth(\mathcal{P}) < d$.

Indeed, let $\mathcal{P} : P = \bigcup_{i=1}^r [F_i, G_i]$ be a partition of P with $sdepth(\mathcal{P}) = sdepth(P)$. Since $\sigma \in P$, it follows that $\sigma \in [F_i, G_i]$ for some i . If $|G_i| \geq d$, then it follows that $\mathcal{P}_{d,\sigma} \neq \emptyset$, since there exist some subsets in the interval $[F_i, G_i]$ of cardinality d which contain σ , a contradiction! Thus, $|G_i| < d$ and therefore $sdepth(\mathcal{P}) < d$.

We set $d = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + 1$ and $\sigma = (a(1), a(2, 1), \dots, a(3r_1, 1), \dots, a(2, k_1), \dots, a(3r_{k_1}, k_1), b(2, 1), \dots, b(3s_1+1, 1), \dots, b(2, k_2), \dots, b(3s_{k_2}+1, k_2), c(2, 1), \dots, c(3t_1+2, 1), \dots, c(2, k_3), \dots, c(3t_{k_3}+2, k_3)) \in \mathbb{N}^n$, where $a(1) = 1$, for any $1 \leq i \leq k_1$, $1 \leq j \leq r_i - 1$, $a(l, i) = \begin{cases} 1 : l = 3j + 1 \text{ or } l = 3r_i \\ 0 : \text{otherwise} \end{cases}$, for any $1 \leq i \leq k_2$, $1 \leq j \leq s_i$, $b(l, i) = \begin{cases} 1 : l = 3j + 1 \\ 0 : \text{otherwise} \end{cases}$, and for any $1 \leq i \leq k_3$, $1 \leq j \leq t_i$, $c(l, i) = \begin{cases} 1 : l = 3j + 1 \\ 0 : \text{otherwise} \end{cases}$. We obtain that $\mathcal{P}_{d+1,\sigma} = \emptyset$. Indeed, if monomial

$$u = x_1 \prod_{i=1}^{k_1} (x_{3r_i, i} \prod_{j=1}^{r_i-1} x_{3j+1, i}) \prod_{i=1}^{k_2} (\prod_{j=1}^{s_i} y_{3j+1, i}) \prod_{i=1}^{k_3} (\prod_{j=1}^{t_i} z_{3j+1, i}),$$

one can easily see that if $a = (a(1), a(2, 1), \dots, a(3r_1, 1), \dots, a(2, k_1), \dots, a(3r_{k_1}, k_1), b(2, 1), \dots, b(3s_1+1, 1), \dots, b(2, k_2), \dots, b(3s_{k_2}+1, k_2), c(2, 1), \dots, c(3t_1+2, 1), \dots, c(2, k_3), \dots, c(3t_{k_3}+2, k_3)) \in \mathbb{N}^n$ such that $a(l, i) \neq 0$ for some $l \notin \{3j+1, 3r_i \mid 1 \leq j \leq r_i - 1, 1 \leq i \leq k_1\}$ or $b(l, i) \neq 0$ for some $l \notin \{3j+1 \mid 1 \leq j \leq s_i, 1 \leq i \leq k_2\}$ or $c(l, i) \neq 0$ for some $l \notin \{3j+1 \mid 1 \leq j \leq t_i, 1 \leq i \leq k_3\}$, then $u \cdot x^a \in I$. Therefore, by previous remark, $sdepth\left(\frac{S}{I}\right) = sdepth(\mathcal{P}) \leq d = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + 1$, as required. \square

4. DEPTH AND STANLEY DEPTH OF THE EDGE IDEALS OF SOME m -CYCLIC GRAPHS WITH A COMMON VERTEX

In this section, we will give some formulas for depth and Stanley depth of the quotient rings of the edge ideals of some m -cyclic graphs with a common vertex. We assume that G is the m -cyclic graph formed by joining m cycles $C_{3r_1}, \dots, C_{3r_{k_1}}, C_{3s_1+1}, \dots, C_{3s_{k_2}+1}, C_{3t_1+2}, \dots, C_{3t_{k_3}+2}$ at a common vertex, where $k_1 + k_2 + k_3 = m$ and $k_i \geq 0$ for $i = 1, 2, 3$. We adopt the following notation to edges of graph G : $E(C_{3r_i, i}) = \{x_1 x_{2, i}, x_{2, i} x_{3, i}, \dots, x_{3r_i, i} x_1\}$ for any $1 \leq i \leq k_1$ and $r_1 \leq r_2 \leq \dots \leq r_{k_1}$, $E(C_{3s_i+1, i}) = \{x_1 y_{2, i}, y_{2, i} y_{3, i}, \dots, y_{3s_i+1, i} x_1\}$ for any $1 \leq i \leq k_2$ and $s_1 \leq \dots \leq s_{k_2}$, $E(C_{3t_i+2, i}) = \{x_1 z_{2, i}, z_{2, i} z_{3, i}, \dots, z_{3t_i+2, i} x_1\}$ for all $1 \leq i \leq k_3$ and $t_1 \leq t_2 \leq \dots \leq t_{k_3}$. Let K be any field, $S = K[x_1, x_{2, 1}, \dots, x_{3r_1, 1}, \dots, x_{2, k_1}, \dots, x_{3r_{k_1}, k_1}, y_{2, 1}, \dots, y_{3s_1+1, 1}, \dots, y_{2, k_2}, \dots, y_{3s_{k_2}+1, k_2}, z_{2, 1}, \dots, z_{3t_1+2, 1}, \dots, z_{2, k_3}, \dots, z_{3t_{k_3}+2, k_3}]$ the polynomial ring. Then the edge ideal of graph G is $J = (x_1 x_{2, 1}, x_{2, 1} x_{3, 1}, \dots, x_{3r_1, 1} x_1, \dots, x_1 x_{2, k_1}, x_{2, k_1} x_{3, k_1}, \dots, x_{3r_{k_1}, k_1} x_1, x_1 y_{2, 1}, y_{2, 1} y_{3, 1}, \dots, y_{3s_1+1, 1} x_1, \dots, x_1 y_{2, k_2}, y_{2, k_2} y_{3, k_2}, \dots, y_{3s_{k_2}+1, k_2} x_1, x_1 z_{2, 1}, z_{2, 1} z_{3, 1}, \dots, z_{3t_1+2, 1} x_1, \dots, x_1 z_{2, k_3}, z_{2, k_3} z_{3, k_3}, \dots, z_{3t_{k_3}+2, k_3} x_1)$.

Example 4.1. The following graph G is the union of 5 circle graphs C_3, C_4, C_5, C_6 and C_7 with a common vertex x_1 .




Figure 2

The edge ideal of graph G is $J = (x_1x_{2,1}, x_{2,1}x_{3,1}, x_{3,1}x_1, x_1x_{2,2}, x_{2,2}x_{3,2}, x_{3,2}x_{4,2}, x_{4,2}x_{5,2}, x_{5,2}x_{6,2}, x_{6,2}x_1, x_1y_{2,1}, y_{2,1}y_{3,1}, y_{3,1}y_{4,1}, y_{4,1}x_1, x_1y_{2,2}, y_{2,2}y_{3,2}, y_{3,2}y_{4,2}, y_{4,2}y_{5,2}, y_{5,2}y_{6,2}, y_{6,2}y_{7,2}, y_{7,2}x_1, x_1z_{2,1}, z_{2,1}z_{3,1}, z_{3,1}z_{4,1}, z_{4,1}z_{5,1}, z_{5,1}x_1)$.

Now, we prove the main results of this section.

Theorem 4.2. *Let G be a graph consisting of the union of cycle graphs $C_{3r_1}, \dots, C_{3r_{k_1}}, C_{3s_1+1}, \dots, C_{3s_{k_2}+1}$ and $C_{3t_1+2}, \dots, C_{3t_{k_3}+2}$ with a common vertex x_1 , where $k_i \geq 0$ for $i = 1, 2, 3$. Let J be its edge ideal. Then*

$$sdepth\left(\frac{S}{J}\right) \geq depth\left(\frac{S}{J}\right) = \begin{cases} \sum_{i=1}^{k_2} s_i, & \text{if } k_1 = k_3 = 0, \\ \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1, & \text{otherwise.} \end{cases}$$

In particular, S/J satisfies the Stanley conjecture.

Proof. Notice that $(J : x_1) = (x_{2,1}, \dots, x_{2,k_1}, x_{3r_1,1}, \dots, x_{3r_{k_1},k_1}, y_{2,1}, \dots, y_{2,k_2}, y_{3s_1+1,1}, \dots, y_{3s_{k_2}+1,k_2}, z_{2,1}, \dots, z_{2,k_3}, z_{3t_1+2,1}, \dots, z_{3t_{k_3}+2,k_3}, x_{3,1}x_{4,1}, \dots, x_{3r_1-2,1}x_{3r_1-1,1}, \dots, x_{3,k_1}x_{4,k_1}, \dots, x_{3r_{k_1}-2,k_1}x_{3r_{k_1}-1,k_1}, y_{3,1}y_{4,1}, \dots, y_{3s_1-1,1}y_{3s_1,1}, \dots, y_{3,k_2}y_{4,k_2}, \dots, y_{3s_{k_2}-1,k_2}y_{3s_{k_2},k_2}, z_{3,1}z_{4,1}, \dots, z_{3t_1,1}z_{3t_1+1,1}, \dots, z_{3,k_3}z_{4,k_3}, \dots, z_{3t_{k_3},k_3}z_{3t_{k_3}+1,k_3})$ and $(J, x_1) = (x_{2,1}x_{3,1}, \dots, x_{3r_1-1,1}x_{3r_1,1}, \dots, x_{2,k_1}x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1}, y_{2,1}y_{3,1}, \dots, y_{3s_1,1}y_{3s_1+1,1}, \dots, y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2}, z_{2,1}z_{3,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1}, \dots, z_{2,k_3}z_{3,k_3}, \dots, z_{3t_{k_3}+1,k_3}z_{3t_{k_3}+2,k_3}, x_1)$, thus we get that

$$\begin{aligned} \frac{S}{(J : x_1)} &\cong \frac{K[x_{3,1}, \dots, x_{3r_1-1,1}]}{(x_{3,1}x_{4,1}, \dots, x_{3r_1-2,1}x_{3r_1-1,1})} \otimes_K \cdots \otimes_K \frac{K[x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1}]}{(x_{3,k_1}x_{4,k_1}, \dots, x_{3r_{k_1}-2,k_1}x_{3r_{k_1}-1,k_1})} \\ &\otimes_K \frac{K[y_{3,1}, \dots, y_{3s_1,1}]}{(y_{3,1}y_{4,1}, \dots, y_{3s_1-1,1}y_{3s_1,1})} \otimes_K \cdots \otimes_K \frac{K[y_{3,k_2}, \dots, y_{3s_{k_2},k_2}]}{(y_{3,k_2}y_{4,k_2}, \dots, y_{3s_{k_2}-1,k_2}y_{3s_{k_2},k_2})} \\ &\otimes_K \frac{K[z_{3,1}, \dots, z_{3t_1+1,1}]}{(z_{3,1}z_{4,1}, \dots, z_{3t_1,1}z_{3t_1+1,1})} \otimes_K \cdots \otimes_K \frac{K[z_{3,k_3}, \dots, z_{3t_{k_3}+1,k_3}]}{(z_{3,k_3}z_{4,k_3}, \dots, z_{3t_{k_3},k_3}z_{3t_{k_3}+1,k_3})} \otimes_K K[x_1], \end{aligned}$$

and

$$\begin{aligned} \frac{S}{(J, x_1)} &\cong \frac{K[x_{2,1}, \dots, x_{3r_1,1}]}{(x_{2,1}x_{3,1}, \dots, x_{3r_1-1,1}x_{3r_1,1})} \otimes_K \cdots \otimes_K \frac{K[x_{2,k_1}, \dots, x_{3r_{k_1},k_1}]}{(x_{2,k_1}x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1})} \\ &\otimes_K \frac{K[y_{2,1}, \dots, y_{3s_1+1,1}]}{(y_{2,1}y_{3,1}, \dots, y_{3s_1,1}y_{3s_1+1,1})} \otimes_K \cdots \otimes_K \frac{K[y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})} \end{aligned}$$

$$\otimes_K \frac{K[z_{2,1}, \dots, z_{3t_1+2,1}]}{(z_{2,1}z_{3,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1})} \otimes_K \cdots \otimes_K \frac{K[z_{2,k_3}, \dots, z_{3t_{k_3}+2,k_3}]}{(z_{2,k_3}z_{3,k_3}, \dots, z_{3t_{k_3}+1,k_3}z_{3t_{k_3}+2,k_3})}.$$

Therefore, by Lemmas 2.6, 2.7, 3.2 and [13, Proposition 2.2.20, Theorem 2.2.21], we obtain that $\text{sdepth}(\frac{S}{J:x_1}) \geq \text{depth}(\frac{S}{J:x_1}) = \sum_{i=1}^{k_1} \lceil \frac{3r_i-3}{3} \rceil + \sum_{i=1}^{k_2} \lceil \frac{3s_i-2}{3} \rceil + \sum_{i=1}^{k_3} \lceil \frac{3t_i-1}{3} \rceil + 1 = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1$,

and $\text{sdepth}(\frac{S}{(J,x_1)}) \geq \text{depth}(\frac{S}{(J,x_1)}) = \sum_{i=1}^{k_1} \lceil \frac{3r_i-1}{3} \rceil + \sum_{i=1}^{k_2} \lceil \frac{3s_i}{3} \rceil + \sum_{i=1}^{k_3} \lceil \frac{3t_i+1}{3} \rceil = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i + k_3$.

Using Lemma 2.5 on the short exact sequence

$$(3) \quad 0 \longrightarrow S/(J : x_1) \xrightarrow{\cdot x_1} S/J \longrightarrow S/(J, x_1) \longrightarrow 0,$$

we conclude that $\text{sdepth}(\frac{S}{J}) \geq \begin{cases} \sum_{i=1}^{k_2} s_i, & \text{if } k_1 = k_3 = 0, \\ \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1, & \text{otherwise.} \end{cases}$

If $k_1 \neq 0$ or $k_3 \neq 0$, then $\text{depth}(\frac{S}{(J,x_1)}) \geq \text{depth}(\frac{S}{J})$. Using Lemma 2.4 on the short exact sequence

$$(3), \text{ it follows that } \text{depth}(\frac{S}{J}) = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1.$$

Assume that $k_1 = k_3 = 0$. We claim that there exists the S -module isomorphism

$$\begin{aligned} \frac{(J:x_1)}{J} &\cong y_{2,1} \left(\frac{K[y_{4,1}, \dots, y_{3s_1+1,1}, y_{2,2}, \dots, y_{3s_2+1,2}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{4,1}y_{5,1}, \dots, y_{3s_1,1}y_{3s_1+1,1}, y_{2,2}y_{3,2}, \dots, y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})} \right) [y_{2,1}] \\ &\oplus y_{3s_1+1,1} \left(\frac{K[y_{3,1}, \dots, y_{3s_1-1,1}, y_{2,2}, \dots, y_{3s_2+1,2}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{3,1}y_{4,1}, \dots, y_{3s_1-2,1}y_{3s_1-1,1}, \dots, y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})} \right) [y_{3s_1+1,1}] \\ &\oplus y_{2,2} \left(\frac{K[y_{3,1}, \dots, y_{3s_1,1}, y_{4,2}, \dots, y_{3s_2+1,2}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{3,1}y_{4,1}, \dots, y_{3s_1-1,1}y_{3s_1,1}, y_{4,2}y_{5,2}, \dots, y_{3s_2,2}y_{3s_2+1,2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})} \right) [y_{2,2}] \\ &\oplus y_{3s_2+1,2} \left(\frac{K[y_{3,1}, \dots, y_{3s_1,1}, y_{3,2}, \dots, y_{3s_2-1,2}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{3,1}y_{4,1}, \dots, y_{3,2}y_{4,2}, \dots, y_{3s_2-2,2}y_{3s_2-1,2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})} \right) [y_{3s_2+1,2}] \\ &\oplus \cdots \\ &\oplus y_{2,k_2} \left(\frac{K[y_{3,1}, \dots, y_{3s_1,1}, \dots, y_{3,k_2-1}, \dots, y_{3s_{k_2-1},k_2-1}, y_{4,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{3,1}y_{4,1}, \dots, y_{3s_1-1,1}y_{3s_1,1}, \dots, y_{4,k_2}y_{5,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})} \right) [y_{2,k_2}] \\ &\oplus y_{3s_{k_2}+1,k_2} \left(\frac{K[y_{3,1}, \dots, y_{3s_1,1}, \dots, y_{3,k_2-1}, \dots, y_{3s_{k_2-1},k_2-1}, y_{3,k_2}, \dots, y_{3s_{k_2}-1,k_2}]}{(y_{3,1}y_{4,1}, \dots, y_{3s_1-1,1}y_{3s_1,1}, \dots, y_{3,k_2}y_{4,k_2}, \dots, y_{3s_{k_2}-2,k_2}y_{3s_{k_2}-1,k_2})} \right) [y_{3s_{k_2}+1,k_2}] \end{aligned}$$

Indeed, if $u \in (J : x_1)$ is a monomial such that $u \notin J$, then there exists some $i \in \{1, \dots, k_2\}$ such that $y_{j,i} \mid u$, where $j = 2$ or $3s_i + 1$.

If $y_{2,1} \mid u$, then we can write u as $u = y_{2,1}^\alpha v$ with $\alpha \geq 1$ and $y_{2,1} \nmid v$. Since $u \notin J$, we have that $v \in K[y_{4,1}, \dots, y_{3s_1+1,1}, y_{2,2}, \dots, y_{3s_2+1,2}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]$ and $v \notin (y_{4,1}y_{5,1}, \dots, y_{3s_1,1}y_{3s_1+1,1}, y_{2,2}y_{3,2}, \dots, y_{3s_2,2}y_{3s_2+1,2}, \dots, y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})$. Similarly, if $y_{3s_1+1,1} \mid u$ and $y_{2,1} \nmid u$, then $u = y_{3s_1+1,1}^\alpha v$ with $\alpha \geq 1$ and $v \in K[y_{3,1}, \dots, y_{3s_1-1,1}, y_{2,2}, \dots, y_{3s_2+1,2}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]$ and $v \notin (y_{3,1}y_{4,1}, \dots, y_{3s_1-2,1}y_{3s_1-1,1}, y_{2,2}y_{3,2}, \dots, y_{3s_2,2}y_{3s_2+1,2}, \dots, y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})$. If u is a monomial such that $y_{2,2} \mid u$, $y_{2,1} \nmid u$ and $y_{3s_1+1,1} \nmid u$, then we have that $u = y_{2,2}^\alpha v$ with $\alpha \geq 1$ and $v \in K[y_{3,1}, \dots, y_{3s_1,1}, y_{4,2}, \dots, y_{3s_2+1,2}, \dots, y_{2,k_2}, \dots, y_{3s_{k_2}+1,k_2}]$ and $v \notin (y_{3,1}y_{4,1}, \dots, y_{3s_1-1,1}y_{3s_1,1}, y_{4,2}y_{5,2}, \dots, y_{3s_2,2}y_{3s_2+1,2}, \dots, y_{2,k_2}y_{3,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2})$. Other cases can be shown in a similar way as the above.

Therefore, by [13, Proposition 2.2.20, Theorem 2.2.21] and Lemma 2.6, it follows that $\text{depth}(\frac{J:x_1}{J}) = \sum_{i=1}^{k_2-1} \lceil \frac{3s_i-2}{3} \rceil + \lceil \frac{3s_{k_2}-3}{3} \rceil + 1 = \sum_{i=1}^{k_2} s_i$.

Now, using Lemma 2.4 on the short exact sequence

$$(4) \quad 0 \longrightarrow (J : x_1)/J \longrightarrow S/J \longrightarrow S/(J : x_1) \longrightarrow 0,$$

this completes the proof. \square

Theorem 4.3. *Let G be a graph as in Theorem 4.2 and J be its edge ideal. Then*

$$s\text{depth}(\frac{S}{J}) \leq \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1.$$

Proof. Let $d = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1$ and $\sigma = (a(1), a(2, 1), \dots, a(3r_1, 1), \dots, a(2, k_1), \dots, a(3r_{k_1}, k_1), b(2, 1), \dots, b(3s_1+1, 1), \dots, b(2, k_2), \dots, b(3s_{k_2}+1, k_2), c(2, 1), \dots, c(3t_1+2, 1), \dots, c(2, k_3), \dots, c(3t_{k_3}+2, k_3)) \in \mathbb{N}^n$, where $a(1) = 1$, for any $1 \leq i \leq k_1$, $1 \leq j \leq r_i - 1$, $a(l, i) = \begin{cases} 1 : l = 3j + 1 \\ 0 : \text{otherwise} \end{cases}$, for any $1 \leq i \leq k_2$, $1 \leq j \leq s_i - 1$, $b(l, i) = \begin{cases} 1 : l = 3j + 1 \text{ or } l = 3s_i \\ 0 : \text{otherwise} \end{cases}$, and for any $1 \leq i \leq k_3$, $1 \leq j \leq t_i$, $c(l, i) = \begin{cases} 1 : l = 3j + 1 \\ 0 : \text{otherwise} \end{cases}$. From the proof of Theorem 3.4, it is enough to prove that $\mathcal{P}_{d+1, \sigma} = \emptyset$. Indeed, if monomial

$$u = x_1 \prod_{i=1}^{k_1} \left(\prod_{j=1}^{r_i-1} x_{3j+1, i} \right) \prod_{i=1}^{k_2} \left(y_{3s_i, i} \prod_{j=1}^{s_i-1} y_{3j+1, i} \right) \prod_{i=1}^{k_3} \left(\prod_{j=1}^{t_i} z_{3j+1, i} \right),$$

one can easily see that if $a = (a(1), a(2, 1), \dots, a(3r_1, 1), \dots, a(2, k_1), \dots, a(3r_{k_1}, k_1), b(2, 1), \dots, b(3s_1+1, 1), \dots, b(2, k_2), \dots, b(3s_{k_2}+1, k_2), c(2, 1), \dots, c(3t_1+2, 1), \dots, c(2, k_3), \dots, c(3t_{k_3}+2, k_3)) \in \mathbb{N}^n$ such that $a(l, i) \neq 0$ for some $l \notin \{3j + 1 \mid 1 \leq j \leq r_i - 1, 1 \leq i \leq k_1\}$ or $b(l, i) \neq 0$ for some $l \notin \{3j + 1, 3s_i \mid 1 \leq j \leq s_i, 1 \leq i \leq k_2\}$ or $c(l, i) \neq 0$ for some $l \notin \{3j + 1 \mid 1 \leq j \leq t_i, 1 \leq i \leq k_3\}$, then $u \cdot x^a \in I$. Therefore $\mathcal{P}_{d+1, \sigma} = \emptyset$, thus we obtain the required result. \square

Theorem 4.4. *Let G be a graph as in Theorem 4.2. Then*

$$s\text{depth}(\frac{J}{I}) \geq \text{depth}(\frac{J}{I}) = \begin{cases} \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_3} t_i - k_1 + 2, & \text{if } k_2 = 0, \\ \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1, & \text{otherwise.} \end{cases}$$

In particular, J/I satisfies the Stanley conjecture.

Proof. We have the S-module isomorphism:

$$\begin{aligned} \frac{J}{I} &\cong \bigoplus_{i=1}^{k_1} x_1 x_{3r_i, i} \left(\frac{K[x_{3,1}, \dots, x_{3r_1-1,1}, \dots, x_{3,i-1}, \dots, x_{3r_{i-1}-1, i-1}, x_{3,i}, \dots, \widehat{x_{3r_{i-1}, i}}, x_{3,i+1}, \dots, x_{3r_{i+1}, i+1}, \dots, x_{3, k_1}, \dots, x_{3r_{k_1}, k_1}]}{(x_{3,1}x_{4,1}, \dots, x_{3r_1-2,1}x_{3r_1-1,1}, \dots, x_{3,i}x_{4,i}, \dots, x_{3r_{i-3}, i}x_{3r_{i-2}, i}, x_{3,i+1}x_{4,i+1}, \dots, x_{3r_{k_1}-1, k_1}x_{3r_{k_1}, k_1})} \right) \\ &\otimes_K \frac{K[y_{3,1}, \dots, y_{3s_1+1,1}, \dots, y_{3,k_2}, \dots, y_{3s_{k_2}+1, k_2}]}{(y_{3,1}y_{4,1}, \dots, y_{3s_1-1,1}y_{3s_1+1,1}, \dots, y_{3s_{k_2}-1, k_2}y_{3s_{k_2}+1, k_2})} \otimes_K \frac{K[z_{3,1}, \dots, z_{3t_1+2,1}, \dots, z_{3, k_3}, \dots, z_{3t_{k_3}+2, k_3}]}{(z_{3,1}z_{4,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1}, \dots, z_{3s_{k_3}+1, k_3}z_{3t_{k_3}+2, k_3})} [x_1, x_{3r_i, i}] \\ &\oplus \left(\bigoplus_{i=1}^{k_2} x_1 y_{3s_i+1, i} \left(\frac{K[y_{3,1}, \dots, y_{3s_1,1}, \dots, y_{3,i-1}, \dots, y_{3s_{i-1}, i-1}, y_{3,i}, \dots, \widehat{y_{3s_i, i}}, y_{3,i+1}, \dots, y_{3s_{i+1}, i+1}, \dots, y_{3, k_2}, \dots, y_{3s_{k_2}+1, k_2}]}{(y_{3,1}y_{4,1}, \dots, y_{3s_1-1,1}y_{3s_1+1,1}, \dots, y_{3,i}y_{4,i}, \dots, y_{3s_{i-2}, i}y_{3s_{i-1}, i}, y_{3,i+1}y_{4,i+1}, \dots, y_{3s_{k_2}-1, k_2}y_{3s_{k_2}+1, k_2})} \right) \otimes_K \right. \end{aligned}$$

$$\begin{aligned} & \frac{K[x_{3,1}, \dots, x_{3r_1-1,1}, \dots, x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1}]}{(x_{3,1}x_{4,1}, \dots, x_{3r_1-2,1}x_{3r_1-1,1}, \dots, x_{3r_{k_1}-2,k_1}x_{3r_{k_1}-1,k_1})} \otimes_K \frac{K[z_{3,1}, \dots, z_{3t_1+2,1}, \dots, z_{3,k_3}, \dots, z_{3t_{k_3}+2,k_3}]}{(z_{3,1}z_{4,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1}, \dots, z_{3s_{k_3}+1,k_3}z_{3t_{k_3}+2,k_3})} (x_1, y_{3s_i+1}, i) \\ & \oplus \left(\bigoplus_{i=1}^{k_3} x_1 z_{3t_i+2}, i \left(\frac{K[z_{3,1}, \dots, z_{3t_1+1,1}, \dots, z_{3,i-1}, \dots, z_{3t_{i-1}+1,i-1}, z_{3,i}, \dots, z_{3t_i+1,i}, z_{3,i+1}, \dots, z_{3t_{i+1}+2,i+1}, \dots, z_{3,k_3}, \dots, z_{3t_{k_3}+2,k_3}]}{(z_{3,1}z_{4,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1}, \dots, z_{3,i}z_{4,i}, \dots, z_{3t_{i-1}+1,i}z_{3t_i,i}, z_{3,i+1}z_{4,i+1}, \dots, z_{3t_{k_3}+1,k_3}z_{3t_{k_3}+2,k_3})} \otimes_K \right. \right. \\ & \left. \left. \frac{K[x_{3,1}, \dots, x_{3r_1-1,1}, \dots, x_{3,k_1}, \dots, x_{3r_{k_1}-1,k_1}]}{(x_{3,1}x_{4,1}, \dots, x_{3r_1-2,1}x_{3r_1-1,1}, \dots, x_{3r_{k_1}-2,k_1}x_{3r_{k_1}-1,k_1})} \otimes_K \frac{K[y_{3,1}, \dots, y_{3s_1+1,1}, \dots, y_{3,k_2}, \dots, y_{3s_{k_2}+1,k_2}]}{(y_{3,1}y_{4,1}, \dots, y_{3s_1-1,1}y_{3s_1,1}, \dots, y_{3s_{k_2}-1,k_2}y_{3s_{k_2},k_2})} \right) (x_1, z_{3t_i+2}, i), \end{aligned}$$

where $x_{i,0} = y_{j,0} = z_{l,0} = 0$ for $3 \leq i \leq k_1$, $3 \leq j \leq k_2$ and $3 \leq l \leq k_3$.

Indeed, let $u \in J$ be a monomial such that $u \notin I$, then $x_1 x_{3r_i,i} | u$ or $x_1 y_{3s_j+1,j} | u$ or $x_1 z_{3t_l+2,l} | u$ for some $1 \leq i \leq k_1$ or $1 \leq j \leq k_2$ or $1 \leq l \leq k_3$.

If $x_1 x_{3r_1,1} | u$, then we can write u as $u = x_1^\alpha x_{3r_1,1}^\beta v$ with $\alpha, \beta \geq 1$, $x_1 \nmid v$ and $x_{3r_1,1} \nmid v$. Since $u \notin I$, we have that $v \in K[x_{3,1}, \dots, x_{3r_1-2,1}, x_{3,2}, \dots, x_{3r_2,2}, \dots, x_{3,k_1}, \dots, x_{3r_{k_1},k_1}, y_{3,1}, \dots, y_{3s_1+1,1}, \dots, y_{3,k_2}, \dots, y_{3s_{k_2}+1,k_2}, z_{3,1}, \dots, z_{3t_1+2,1}, \dots, z_{3,k_3}, \dots, z_{3t_{k_3}+2,k_3}]$ and $v \notin (x_{3,1}x_{4,1}, \dots, x_{3r_1-3,1}x_{3r_1-2,1}, x_{3,2}x_{4,2}, \dots, x_{3r_2-1,2}x_{3r_2,2}, \dots, x_{3,k_1}x_{4,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1}, y_{3,1}y_{4,1}, \dots, y_{3s_1,1}y_{3s_1+1,1}, \dots, y_{3,k_2}y_{4,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2}, z_{3,1}z_{4,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1}, \dots, z_{3,k_3}z_{4,k_3}, \dots, z_{3t_{k_3}+1,k_3}z_{3t_{k_3}+2,k_3})$. Similarly, if $x_1 x_{3r_2,2} | u$ and $x_1 x_{3r_1,1} \nmid u$, then $u = x_1^\alpha x_{3r_2,2}^\beta v$ with $\alpha, \beta \geq 1$ and $v \in K[x_{3,1}, \dots, x_{3r_1-1,1}, x_{3,2}, \dots, x_{3r_2-2,2}, x_{3,3}, \dots, x_{3r_3,3}, \dots, x_{3,k_1}, \dots, x_{3r_{k_1},k_1}, y_{3,1}, \dots, y_{3s_1+1,1}, \dots, y_{3,k_2}, \dots, y_{3s_{k_2}+1,k_2}, z_{3,1}, \dots, z_{3t_1+2,1}, \dots, z_{3,k_3}, \dots, z_{3t_{k_3}+2,k_3}]$ and $v \notin (x_{3,1}x_{4,1}, \dots, x_{3r_1-2,1}x_{3r_1-1,1}, x_{3,2}x_{4,2}, \dots, x_{3r_2-3,2}x_{3r_2-2,2}, x_{3,3}x_{4,3}, \dots, x_{3r_3-1,3}x_{3r_3,3}, \dots, x_{3,k_1}x_{4,k_1}, \dots, x_{3r_{k_1}-1,k_1}x_{3r_{k_1},k_1}, y_{3,1}y_{4,1}, \dots, y_{3s_1,1}y_{3s_1+1,1}, \dots, y_{3,k_2}y_{4,k_2}, \dots, y_{3s_{k_2},k_2}y_{3s_{k_2}+1,k_2}, z_{3,1}z_{4,1}, \dots, z_{3t_1+1,1}z_{3t_1+2,1}, \dots, z_{3,k_3}z_{4,k_3}, \dots, z_{3t_{k_3}+1,k_3}z_{3t_{k_3}+2,k_3})$. Other cases can be shown in a similar way as the above.

Therefore, by Lemmas 2.4–2.7, 3.2 and [13, Proposition 2.2.20, Theorem 2.2.21], if $k_2 \neq 0$, then $\text{sdepth}(\frac{J}{I}) \geq \text{depth}(\frac{J}{I}) = \min\left\{\sum_{i=1}^{k_1-1} \lceil \frac{3r_i-3}{3} \rceil + \lceil \frac{3r_{k_1}-4}{3} \rceil + \sum_{i=1}^{k_2} \lceil \frac{3s_i-1}{3} \rceil + \sum_{i=1}^{k_3} \lceil \frac{3t_i}{3} \rceil, \sum_{i=1}^{k_1} \lceil \frac{3r_i-3}{3} \rceil + \sum_{i=1}^{k_2-1} \lceil \frac{3s_i-2}{3} \rceil + \lceil \frac{3s_{k_2}-3}{3} \rceil + \sum_{i=1}^{k_3} \lceil \frac{3t_i}{3} \rceil, \sum_{i=1}^{k_1} \lceil \frac{3r_i-3}{3} \rceil + \sum_{i=1}^{k_2} \lceil \frac{3s_i-2}{3} \rceil + \sum_{i=1}^{k_3-1} \lceil \frac{3t_i-1}{3} \rceil + \lceil \frac{3t_{k_3}-2}{3} \rceil\right\} + 2 = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_2} s_i + \sum_{i=1}^{k_3} t_i - k_1 + 1.$

If $k_2 = 0$, then $\text{sdepth}(\frac{J}{I}) \geq \text{depth}(\frac{J}{I}) = \min\left\{\sum_{i=1}^{k_1-1} \lceil \frac{3r_i-3}{3} \rceil + \lceil \frac{3r_{k_1}-4}{3} \rceil + \sum_{i=1}^{k_3} \lceil \frac{3t_i}{3} \rceil, \sum_{i=1}^{k_1} \lceil \frac{3r_i-3}{3} \rceil + \sum_{i=1}^{k_3-1} \lceil \frac{3t_i-1}{3} \rceil + \lceil \frac{3t_{k_3}-2}{3} \rceil\right\} + 2 = \sum_{i=1}^{k_1} r_i + \sum_{i=1}^{k_3} t_i - k_1 + 2$. This completes the proof. \square

Acknowledgments

The author would like to express her sincere thanks to the editor for help and encouragement. Special thanks are due to the referee for suggesting several improvements to the proof of Theorems 3.4 and 4.3.

REFERENCES

- [1] C. Biró, D. M. Howard, M. T. Keller, W. T. Trotter and S.J. Young, Interval partitions and Stanley depth, *J. Combin. Theory*, Ser. A, **117** (2010) 475-482.
- [2] M. Cimpoeaş, On the Stanley depth of edge ideals of line and cyclic graphs, *Romanian Journal of Math. and Computer Science*. **5** (1), (2015), 70-75.
- [3] CoCoATeam, *CoCoA: a system for doing Computations in Commutative Algebra*, Available at <http://cocoa.dima.unige.it>
- [4] A. M. Duval, B. Goeckneker, C. J. Klivans and J. L. Martine, A non-partitionable Cohen- Macaulay simplicial complex, *arXiv:1504.04279v2*.
- [5] J. Herzog, A Survey on Stanley Depth, *Lecture Notes in Mathematics* 2083, Springer (2013), 3-45.
- [6] J. Herzog, M. Vlădoiu and X. Zheng, How to compute the Stanley depth of a monomial ideal, *J. Algebra* **322** (2009), 3151-3169.

- [7] S. Morey, Depths of powers of the edge ideal of a tree, *Comm. Algebra* **38** (11), (2010), 4042-4055.
- [8] A. Rauf, Depth and sdepth of multigraded module, *Comm. Algebra* **38** (2), (2010), 773-784.
- [9] G. Rinaldo, An algorithm to compute the Stanley depth of monomials ideals, *Le Matematiche*, Vol LXIII (ii) (2008), 243-256.
- [10] R. P. Stanley, Linear Diophantine equations and local cohomology, *Invent. Math.* **68** (2), (1982), 175-193.
- [11] A. Ştefan, Stanley depth of powers of the path ideal, *arXiv:1409.6072V1*.
- [12] W. V. Vasconcelos, *Arithmetic of Blowup Algebras*, London Math. Soc., Lecture Note Series 195. Cambridge: Cambridge University Press, 1994.
- [13] R. H. Villarreal, *Monomial algebras*, Monographs and Textbooks in Pure and Applied Mathematics, 238. Marcel Dekker, Inc., New York, 2001.
- [14] G. J. Zhu, F. Shi and Y. Gu, Arithmetical rank of the edge ideals of some n -cyclic graphs with a common edge, *Turkish Journal of Mathematics* **39** (1), (2015), 112-123.

SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU 215006, P. R. CHINA
E-mail address: zhuguangjun@suda.edu.cn